



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number: **0 675 121 A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **95302076.5**

(51) Int. Cl.⁶: **C07D 333/56, C07D 333/58**

(22) Date of filing: **28.03.95**

(30) Priority: **31.03.94 US 220853**

(43) Date of publication of application:
04.10.95 Bulletin 95/40

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT SE


(71) Applicant: **ELI LILLY AND COMPANY**
Lilly Corporate Center
Indianapolis Indiana 46285 (US)

(72) Inventor: **Dodge, Jeffrey Alan**
11134 Indian Lake Boulevard
Indianapolis, Indiana 46236 (US)
Inventor: **Stocksdale, Mark Gregory**
1257 Long Fellow Avenue
South Bend, Indiana 46615 (US)

(74) Representative: **Hudson, Christopher Mark et al**
Lilly Industries Limited
European Patent Operations
Erl Wood Manor
Windlesham Surrey GU20 6PH (GB)

(54) **Novel intermediates and processes for preparing benzothioephene compounds.**


(57) The present invention provides a novel process for preparing a compound of formula I




wherein

R¹ and R² combine to form C₄-C₈ polymethylene, -CH₂CH(CH₃)CH₂CH₂-, -CH₂C(CH₃)₂CH₂CH₂-, or -CH₂CH₂OCH₂CH₂-; or a pharmaceutically acceptable salt or solvate thereof, comprising

a) reacting a compound of formula II



with a compound of formula III




wherein R¹ and R² are as defined above, in the presence of triphenylphosphine and diethyl azodicarboxylate; or


b) reacting a compound of formula II

EP 0 675 121 A1

ATTORNEY DOCKET NUMBER: 10177-211-999
SERIAL NUMBER: 09/910,388
REFERENCE: B74





with a compound of formula $Z-CH_2-CH_2-Z$ in which Z is the same or different leaving group, and reacting the resulting compound of formula IV



wherein Z is a leaving group, with pyrrolidine, piperidine, hexamethyleneimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine ;

c) deprotecting the 6- and 4'-position hydroxy group of the reaction product of step a) or step b) ; and
d) optionally salifying or forming a solvate of the reaction product of step c).


Also provided are novel intermediates of formulae II and IV



wherein Z is a leaving group.

This invention relates to the fields of pharmaceutical and organic chemistry and provides novel intermediates and processes for preparing certain benzothiophene compounds useful, *inter alia*, for the treatment of osteoporosis in post-menopausal women.


Compounds of formula I




wherein

R¹ and R² combine to form C₄-C₈ polymethylene, -CH₂CH(CH₃)CH₂CH₂-, -CH₂C(CH₃)₂CH₂CH₂-, or -CH₂CH₂OCH₂CH₂-; or a pharmaceutically acceptable salt or solvate thereof, are known as antifertility agents (see, e.g., U.S. Pat. No. 4,133,814) as well as agents which are useful for the treatment of postmenopausal osteoporosis (see, e.g., Draper, *et al.*, *Effects of Raloxifene (LY139481 HCl) on Biochemical Markers of Bone and Lipid Metabolism in Healthy Postmenopausal Women*, Hong Kong, Fourth Int'l. Symp. on Osteoporosis, March 29, 1993). Raloxifene is a compound of formula I in which R¹ and R² are combined to form C₅ polymethylene.

Jones and Suarez, in U.S. Pat. No. 4,133,814, *supra*, first taught the compounds prepared by the present invention, and showed a number of processes for preparing them. In general, Jones, *et al.*, as well as Peters, M.K. (U.S. Pat. No. 4,380,635), prepare benzothiophenes of formula I via the acylation of a compound of formula V



with a compound of formula VI




wherein R³ is chloro or bromo and R¹ and R² are as defined above, and the 6- and 4'-position hydroxy protecting groups are removed via standard procedures.

One aspect of the present invention provides novel processes for preparing compounds of formula I.

Another aspect of the present invention provides novel intermediates which are useful for preparing compounds of formula I.

The present invention provides a process for preparing a compound of formula I




I

wherein


R¹ and R² combine to form C₄-C₈ polymethylene, -CH₂CH(CH₃)CH₂CH₂-, -CH₂C(CH₃)₂CH₂CH₂-, or -CH₂CH₂OCH₂CH₂-; or a pharmaceutically acceptable salt or solvate thereof, comprising

a) reacting a compound of formula II



II


with a compound of formula III



III

wherein R¹ and R² are as defined above, in the presence of triphenylphosphine and diethyl azodicarboxylate; or


b) reacting a compound of formula II



II

with a compound of the formula Z-CH₂-CH₂-Z in which Z is the same or different leaving group, and reacting


the resulting compound of formula IV




wherein Z is a leaving group, with pyrrolidine, piperidine, hexamethyleneimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine;

c) deprotecting the 6- and 4'-position hydroxy group of the reaction product of step a) or step b); and
d) optionally salifying or forming a solvate of the reaction product of step c).

Also provided by the present invention is a compound of formula II



and compounds of formula IV



wherein Z is a leaving group.

The present invention provides a convenient process for preparing benzothiophenes of formula I.

The starting material for the process of the present invention, a compound of formula II, is prepared via known procedures using known or commercially available materials. In general, a compound of formula V, *supra*, which is described by Peters, M.K. in U.S. Pat. No. 4,380,635, is acylated with anisoyl chloride in the presence of an appropriate catalyst and an inert solvent. U.S. Pat. No. 4,380,635 is herein incorporated by reference.

The acylation step of the present process, in which a compound of formula VI is prepared (*see* Scheme I, *infra*), is a Friedel-Crafts reaction, and is carried out in the usual manner, using an appropriate catalyst such

as, for example, aluminum chloride or bromide, gallium dichloride or trichloride, molybdenum pentachloride, iron chloride, and the like [see, e.g., Olah, G.A., *et al.*, *J. Am. Chem. Soc.*, 94:7448 (1974)], in the presence of an inert solvent. Of these catalysts, aluminum chloride is preferred.

Appropriate inert solvents include, for example, halogenated solvents such as dichloromethane, 1,2-dichloroethane, chloroform, and the like, as well as aromatics such as benzene, chlorobenzene, and the like. Of these, a halogenated solvent, especially 1,2-dichloroethane, is preferred.

It has been found that toluene is rather easily acylated under the conditions used in the Friedel-Crafts reaction. It, therefore, is important to remove any previously used toluene so that waste of the acylating agent is avoided.

The above acylation may be carried out at temperatures from about -30° C to about 100° C, preferably in the range from about -10° C to about 20° C.

It generally is preferred that an equimolar amount of each substrate is reacted together. However, it is preferred that an excess of anisoyl chloride is used for this reaction.


The acylation step is rapid, taking from about 15 minutes to a few hours when run at the preferred temperature range. Longer reaction times may be used, but usually are not advantageous.

Next, the methyl hydroxy protecting group of the benzoyl moiety of a formula VI compound is selectively removed via standard procedures. Typically, the formula VI compound is reacted with the lithium or, preferably, the sodium salt of ethanethiol, essentially as described in Preparation 2, *infra*.


This reaction is performed in a non-reactive solvent such as N,N-dimethylformamide and the like. The reaction is performed at temperatures from about 50° C to about 120° C and allowed to run until compounds of formula II are prepared. The reaction typically takes about 4 hours when run at 80° C. However, the progress of the reaction may be monitored by using standard chromatographic techniques.

The formula II compound is novel and useful as the starting material for the processes of the present invention.


Scheme I



V



VI




II


50

The process of the present invention is represented below in Scheme II.

55

Scheme II





III

wherein R^1 and R^2 are as defined above, in the presence of triphenylphosphine, diethyl azodicarboxylate (DEAD), and an appropriate solvent, resulting in a compound of formula VII above. This process is known in the art as a Mitsunobo coupling.

Although this coupling can be accomplished using various concentrations of the reactants and reagents, it is best to use 1 to 2 equivalents of a formula III compound, triphenylphosphine, and DEAD per each equivalent of formula II compound used.

This reaction also is best carried out in the presence of an inert solvent such as, for example, toluene, benzene, or, preferably tetrahydrofuran. The reaction is performed at temperatures from about 0°C to about 40°C , preferably at ambient temperature, until a formula VII compound is prepared. Typically, the reaction takes about 18 hours when run at ambient temperature, but the progress of the reaction can be monitored via standard chromatographic techniques.

A preferred formula III reactant, and thus, providing a compound of formula VII, is that in which R^1 and R^2 combine to form C_4 - C_6 polymethylene, particularly C_6 polymethylene. The term " C_4 - C_6 polymethylene" refers to tetramethylene, pentamethylene, and hexamethylene.

Following preparation of a formula VII compound, the 6- and 4'-position protected hydroxy groups are demethylated via procedures known by one of ordinary skill in the art.

Generally, deprotection (demethylation) is accomplished by the addition of a sulfur compound and a Lewis acid, preferably aluminum chloride, to the reaction mixture remaining from the preparation of formula VII compounds.

The sulfur compound, preferably, are the alkylthiols such as methanethiol, isopropanethiol, butanethiol, ethanethiol, the preferred agent, and the like, dialkyl sulfides such as, for example, butyl *s*-butyl sulfide, ethyl propyl sulfide, butyl isopropyl sulfide, dimethyl sulfide, methyl ethyl sulfide, and the like, benzenethiol, methionine, and alkyl phenyl sulfides such as, for example, methyl phenyl sulfide, ethyl phenyl sulfide, butyl phenyl sulfide, and the like.

It has been found that the demethylation step is best carried out when a substantial excess of the sulfur compound and the Lewis acid is used, in the range from about 4 to about 10 moles per mole of substrate. The process can be carried out, although less efficiently, with a smaller amount of the sulfur compound, in the range from about 2 to 3 moles per mole of substrate. It is also possible to use a small amount of the sulfur compound, such as 2 to 3 moles per mole of substrate, and to improve the yield by the addition of about 1 to 3 moles of an alkali metal halide such as sodium, potassium, or lithium chloride, iodide, or bromide.

This demethylation reaction runs well, and is preferred, when run at ambient temperature, in the range from about 15°C to about 30°C . However, the demethylation step may be carried out at temperatures in the range from about -30°C to about 50°C . Short reaction times, about 1 hour, have been found to be adequate.

After a compound of formula VII has been demethylated, forming a compound of formula I, it is recovered and isolated by conventional means. The examples below further illustrate such recovery and isolation.

The above-described process can be carried out in two, isolated steps in which the product of the first step is recovered, isolated, and purified prior to carrying out the demethylation step. However, the present process also may be carried out in the same vessel without purification of intermediates, thus providing a one-pot process.

As an alternative to the above-described process, a formula II compound is first alkylated with a compound of the formula $\text{Z}-\text{CH}_2-\text{CH}_2-\text{Z}$, in which each Z is the same or different leaving group, in the presence of an alkali solution, and reacting the product of the first step with pyrrolidine, piperidine, hexamethylenimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine. The reaction product from this amine addition step is then demethylated to provide a compound of formula I (see, Scheme II, *supra*).

First, the phenol moiety of the formula II compound is converted to an ether moiety via procedures known in the art. Typically, an excess of the alkylating agent ($\text{Z}-\text{CH}_2-\text{CH}_2-\text{Z}$) is reacted with the formula II compound in an alkali solution.

Appropriate alkylating agents, in which each Z is the same or different leaving group, include, for example, the sulfonates such as methanesulfonate, 4-bromobenzenesulfonate, toluenesulfonate, ethanesulfonate, isopropanesulfonate, 4-methoxybenzenesulfonate, 4-nitrobenzenesulfonate, 2-chlorobenzenesulfonate, triflate, and the like, halogens such as bromo, chloro, iodo, and the like, and other related leaving groups. Because

bromo is preferred, the preferred alkylating agent is 1,2-dibromoethane. This alkylation is carried out by using at least 2 equivalents, and preferably greater than 2 equivalents, of alkylating agent per equivalent of substrate.

In an alkali solution, the phenol moiety exists as a phenoxide ion which displaces one of the leaving groups of the alkylating agents. A preferred alkali solution contains potassium carbonate in an inert solvent such as, for example, methylethyl ketone or dimethylformamide.

This reaction is best run when the alkali solution containing the reactants and reagents is brought to reflux and allowed to run to completion. When using methylethyl ketone as the preferred solvent, reaction times run from about 6 hours to about 20 hours.

The reaction product from the first step, a novel compound of formula IV, is then reacted with pyrrolidine, piperidine, hexamethylenimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine, forming a compound of VII, via standard techniques. Preferably, the hydrochloride salt of piperidine is reacted with the formula IV compound in an inert solvent, such as anhydrous dimethylformamide, and heated to a temperature range from about 60° C to about 110° C. When the mixture is heated to a preferred temperature of about 90° C, the reaction only takes about 30 minutes to about 1 hour. However, changes in the reaction conditions will influence the amount of time this reaction needs to be run to completion. Of course, the progress of the reaction can be monitored via standard chromatographic techniques.

Once this reaction is completed, the resulting formula VII compound is demethylated according to the above-described process, forming a compound of formula I.

Although the reaction product from each step of the present process can be recovered and isolated via standard techniques, each step of this process can be carried out in a single vessel, providing a one-pot process.

Although the free-base form of formula I compounds can be used for various medicinal indications, it is preferred to prepare and use a pharmaceutically acceptable salt form. Thus, compounds of formula I optionally form pharmaceutically acceptable acid and base addition salts with a wide variety of organic and inorganic acids and bases, and include the physiologically acceptable salts which are often used in pharmaceutical chemistry. Preparation of such salts are also part of this invention. Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, phosphoric, hypophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, β -hydroxybutyrate, butyne-1,4-dioate, hexyne-1,4-dioate, caprate, caprylate, chloride, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, terephthalate, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, sulfate, bisulfate, pyrosulfate, sulfite, bisulfite, sulfonate, benzenesulfonate, p-bromophenylsulfonate, chlorobenzene sulfonate, ethanesulfonate, 2-hydroxyethanesulfonate, methanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, p-toluenesulfonate, xylenesulfonate, tartarate, and the like. A preferred salt is the hydrochloride salt.


The pharmaceutically acceptable acid addition salts are typically formed by reacting a compound of formula I with an equimolar or excess amount of acid. The reactants are generally combined in a mutual solvent such as diethyl ether or ethyl acetate. The salt normally precipitates out of solution within about one hour to 10 days and can be isolated by filtration or the solvent can be stripped off by conventional means.

Bases commonly used for formation of salts include ammonium hydroxide and alkali and alkaline earth metal hydroxides, carbonates, as well as aliphatic and primary, secondary and tertiary amines, aliphatic diamines. Bases especially useful in the preparation of addition salts include ammonium hydroxide, potassium carbonate, methylamine, diethylamine, ethylene diamine and cyclohexylamine.


The pharmaceutically acceptable salts and solvates, the latter of which are prepared by methods well known to one of ordinary skill in the organic chemical arts, generally have enhanced solubility characteristics compared to the compound from which they are derived, and thus are often more amenable to formulation as liquids or emulsions.

The following examples are presented to further illustrate the preparation of compounds of the present invention. It is not intended that the invention be limited in scope by reason of any of the following examples.

The terms "NMR", "IR" or "MS" following a synthesis protocol indicates that the nuclear magnetic resonance spectrum, infrared spectrum, or the mass spectrometry was performed and was consistent with the title product.

Preparation 1**6-methoxy-2-(*p*-methoxyphenyl)benzo[*b*]thien-3-yl *p*-methoxyphenyl ketone**

To a solution of *p*-anisoyl chloride (1.54 g, 9.00 mmol) stirring at ambient temperature in anhydrous methylene chloride (100 ml) was added 6-methoxy-2-(4-methoxyphenyl)benzo[*b*]thiophene (1.62 g, 6.00 mmol). The resulting mixture was cooled to 0° C and aluminum chloride (1.20 g, 9.00 mmol) was added in small portions over a five minute period. After 1 hour, the reaction mixture was poured into ice water (150 mL) and extracted with methylene chloride. The combined organic extracts were washed with 1N sodium hydroxide, water, brine, dried (magnesium sulfate), filtered, and concentrated. The resulting material was purified by flash chromatography (silica gel, 30% ethyl acetate in hexanes) giving 2.25 g (93%) of the desired product as a light yellow solid. The product was further purified by recrystallization from acetone/methanol to give 2.11 g (87%): ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.64-7.69 (m, 3H), 7.29-7.32 (m, 3H), 6.86-7.00 (m, 5H), 3.83 (s, 3H) 3.76 (s, 3H); ¹³C NMR (75.5 MHz, DMSO-*d*₆) δ 192.0, 163.6, 159.5, 157.4, 141.0, 139.4, 133.2, 131.8, 130.0, 129.6, 125.2, 123.3, 115.0, 114.4, 114.1, 105.1, 55.5, 55.1; IR (CHCl₃) 3020, 3015, 2970, 2940, 2840, 1600, 1475, 1253, 1218, 1167; MS (FD) 404 (M⁺); Anal. Calcd for C₂₄H₂₀O₄S: C, 71.27; H, 4.98; S, 7.93; O, 15.82. Found: C, 71.50; H, 5.00; S, 7.98; O, 15.77.


Example 1**(4-hydroxyphenyl) [6-methoxy-2-(4-methoxyphenyl)benzo[*b*]thien-3-yl]methano**

To a solution of 6-methoxy-2-(*p*-methoxyphenyl) benzo[*b*]thien-3-yl *p*-methoxyphenyl ketone (0.40 g, 1.00 mmol) stirring in dimethylformamide (DMF; 2 mL) at ambient temperature was added the sodium salt of ethanethiol (3.0 ml of 0.50 M solution in DMF) and the reaction mixture was heated to 80° C. After 4 hours, the reaction was diluted with ethyl acetate, and water was added. The mixture was then neutralized with 1N hydrochloric acid and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried (magnesium sulfate), filtered, and concentrated. The resulting solid was purified by radial chromatography (silica gel, 2 mm, 5% EtOAc in methylene chloride) to give 0.31 g (79 %) of the desired product as a yellow foam: ¹H-NMR (300 MHz, CDCl₃) δ 7.70-7.73 (d, 2H, J = 8.6 Hz); 7.53 (d, 1H, J = 8.5 Hz), 7.31-7.34 (m, 3H), 6.94-6.98 (dd, 1H, J = 9.0 Hz, J = 2.4 Hz), 6.75 (d, 2H, J = 8.7 Hz), 6.67 (d, 2H, J = 9.1 Hz), 3.88 (s, 3H), 3.74 (s,

3H); ^{13}C NMR (75.5 MHz CDCl_3) δ 192.9, 159.9, 158.6, 156.5, 141.9, 138.9, 132.7, 131.7, 129.2, 129.1, 128.8, 124.7, 122.8, 114.3, 113.7, 112.9, 103.4, 54.5, 54.1; IR (CHCl_3) 3585, 3265, 3022, 3012, 2970, 2940, 2840, 1602, 1476, 1254, 1163; MS (FD+) 390 (M+); Anal. Calcd for $\text{C}_{23}\text{H}_{18}\text{O}_4\text{S}$: C, 70.75; H, 4.65. Found: C, 70.93; H, 4.56.

Example 2

[6-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl][4-[2-(1-piperidinyl)ethoxy]phenyl]methanone




VII

To a solution of 1-piperidineethanol (0.32 g, 2.50 mmol) stirring in anhydrous tetrahydrofuran (20 mL) at ambient temperature was added triphenylphosphine (0.52 g, 2.00 mmol) followed by diethyl azodicarboxylate (2.0 mmol). After 5 minutes, (4-hydroxyphenyl)[b-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl]methano (0.39 g, 1.00 mmol) was added to this solution and the mixture stirred for 18 hours at ambient temperature. The reaction mixture was concentrated and the resulting material purified by radial chromatography (silica gel, 4/1/1 methylene chloride/ethyl acetate/methanol) to give 0.46 g (91 %) the desired product as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 7.76 (d, 2H, J = 8.8 Hz), 7.52 (d, 1H, J = 8.9 Hz), 7.31-7.35 (m, 3H), 6.95 (dd, 1H, J = 8.9, 2.3 Hz), 6.74-6.77 (m, 4H), 4.08 (t, 2H, J = 6.6 Hz), 3.88 (s, 3H), 3.75 (s, 3H), 2.73 (t, 2H, J = 6.6 Hz), 2.44-2.52 (m, 4H), 1.55-1.62 (m, 4H), 1.39-1.46 (m, 2H); ^{13}C NMR (75.5 MHz CDCl_3) δ 193.2, 163.0, 159.7, 157.6, 142.5, 140.1, 132.3, 130.6, 130.4, 130.3, 126.0, 124.1, 114.8, 114.2, 114.1, 104.5, 66.3, 57.7, 55.6, 55.2, 55.1, 25.9, 24.1; IR (CHCl_3) 3008, 2941, 2854, 2833, 1646, 1599, 1476, 1254, 1167 cm^{-1} ; MS (FD) 501 (M+).

Example 3

[6-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl][4-[2-(1-bromo)ethoxy]phenyl]methanone




IV

To a solution of (4-hydroxyphenyl) [b-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl]methano (3.90 g, 10.0 mmol) stirring in methylethyl ketone (25 mL) was added ground potassium carbonate (2.07 g, 15.0 mmol) followed by 1,2-dibromoethane (10 mL) and the solution brought to reflux. After 18 hours, the reaction mixture was hot filtered and the residue washed thoroughly with ethyl acetate. The filtrate was concentrated and the resulting material purified by flash column chromatography (silica gel, 50% ethyl acetate/hexanes) to give 4.32 g (87 %) of desired product as a yellow solid: ^1H -NMR (300 MHz, CDCl_3) δ 7.76 (d, 2H, J = 8.8 Hz), 7.55 (d, 1H, 8.9 Hz), 7.31-7.35 (m, 3H), 6.96 (dd, 1H, J = 8.9 Hz, J = 2.3 Hz), 6.74-6.78 (m, 4H); IR (CHCl_3) 3030, 3015,

2965, 2942, 2835, 1601, 1475, 1253, 1240, 1167 cm^{-1} ; MS (FD+) 496 (M^+Br^{79}), 498 (M^+Br^{81}); Anal. calcd. for $\text{C}_{26}\text{H}_{21}\text{BrO}_4\text{S}$: C, 60.37; H, 4.26; Br, 16.07. Found: C, 60.22; H, 4.54; Br, 16.20.

Example 4

[6-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl][4-[2-(1-piperidinyl)ethoxy]phenyl]methanone




VII

To [6-methoxy-2-(4-methoxyphenyl)benzo[b]thien-3-yl][4-[2-(1-bromo)ethoxy]phenyl]methanone (0.497 g, 1.00 mmol) stirring at ambient temperature in anhydrous DMF (10 mL) was added piperidine (3.00 mmol). The mixture was heated to 90° C. After 45 minutes, the solution was cooled to ambient temperature and water was added. The reaction mixture was extracted with ethyl acetate and the combined organic extracts washed with brine, dried (magnesium sulfate), filtered, and concentrated. The resulting material was purified by radial chromatography (silica gel, 10% MeOH in ethyl acetate) to give 0.47 g (93%) the desired product as a yellow solid: ^1H NMR (300 MHz, CDCl_3) δ 7.75 (d, 2H, J = 8.7 Hz), 7.52 (d, 1H, J = 8.9 Hz), 7.31-7.35 (m, 3H), 6.95 (dd, 1H, J = 8.9, 2.3 Hz), 6.74-6.77 (m, 4H), 4.08 (t, 2H, J = 6.6 Hz), 3.88 (s, 3H), 3.75 (s, 3H), 2.73 (t, 2H, J = 6.6 Hz), 2.44-2.52 (m, 4H), 1.55-1.62 (m, 4H), 1.39-1.46 (m, 2H); ^{13}C NMR (75.5 MHz CDCl_3) δ 193.2, 163.1, 160.0, 157.7, 142.2, 140.1, 133.9, 132.3, 130.6, 130.4, 130.3, 126.0, 124.1, 114.8, 114.2, 114.1, 104.5, 66.3, 57.7, 55.6, 55.2, 55.1, 25.9, 24.1; IR (CHCl_3) 3008, 2941, 2854, 2834, 1646, 1599, 1476, 1254, 1167 cm^{-1} ; MS (FD) 502 (M^+).

Example 5

[6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thien-3-yl][4-[2-(1-piperidinyl)ethoxy]phenyl]methanone hydrochloride



I


To a 250 ml three-neck round bottom flask are combined 50 mL of ethylene chloride, aluminum trichloride (9.6 g, 72 mmol) and ethanethiol (6.39 g, 103 mmol) to create a pale yellow liquid. To this liquid is added the product of Example 4 (5.0 g, 10.3 mmol) in a gradual fashion. An oil precipitate and the mixture is stirred for about 20 minutes. After cooling the reaction mixture in an ice bath, 100 mL of tetrahydrofuran is added and the mixture is allowed to stir until all of the oil goes into solution.

The reaction mixture is then poured over ice (200 mL) and water (500 mL) and concentrated hydrochloric acid (10 mL) is added. The resulting precipitate (oil) is separated from the liquid by decantation. The liquid is extracted with warm chloroform (2 x 300 mL). The oil is dissolved by mixing with ethyl acetate, chloroform,

sodium bicarbonate, and a small amount of sodium hydroxide. The chloroform extract and the dissolved oil are transferred to a separatory funnel and washed with sodium bicarbonate. The organic phase is then dried over magnesium sulfate and the solvents are removed by evaporation. The resulting product is further purified by high performance liquid chromatography.

Claims

1. A process for preparing a compound of formula I




I

wherein


R¹ and R² combine to form C₄-C₈ polymethylene, -CH₂CH(CH₃)CH₂CH₂-, -CH₂C(CH₃)₂CH₂CH₂-, or -CH₂CH₂OCH₂CH₂-; or a pharmaceutically acceptable salt or solvate thereof, comprising

a) reacting a compound of formula II



II


with a compound of formula III




III

wherein R¹ and R² are as defined above, in the presence of triphenylphosphine and diethyl azodicarboxylate; or

b) reacting a compound of formula II




15 with a compound of formula $Z-CH_2-CH_2-Z$ in which Z is the same or different leaving group, and reacting the resulting compound of formula IV



30 wherein Z is a leaving group, with pyrrolidine, piperidine, hexamethyleneimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine;

c) deprotecting the 6- and 4'-position hydroxy group of the reaction product of step a) or step b); and
d) optionally salifying or forming a solvate of the reaction product of step c).


- 35 2. A process according to Claim 1 wherein R^1 and R^2 combine to form C_4 polymethylene and the salt thereof is the hydrochloride salt.
3. A process according to any one of Claims 1 to 2 wherein steps a), c), and d) are used.
- 40 4. A process according to any one of Claims 1 to 2 wherein steps b), c), and d) are used.
5. A process according to Claim 4 wherein Z of said formula IV compound is bromo.
6. A process according to any one of Claims 1 to 5 wherein the process is carried out in a single vessel.
- 45 7. A compound of formula I




wherein

R¹ and R² combine to form C₄-C₆ polymethylene, -CH₂CH(CH₃)CH₂CH₂-, -CH₂C(CH₃)₂CH₂CH₂-, or -CH₂CH₂OCH₂CH₂-; or a pharmaceutically acceptable salt or solvate thereof, when prepared by the process comprising

a) reacting a compound of formula II



with a compound of formula III




wherein R¹ and R² are as defined above, in the presence of triphenylphosphine and diethyl azodicarboxylate; or

b) reacting a compound of formula II

5

10




15

with a compound of formula $Z-CH_2-CH_2-Z$ in which Z is the same or different leaving group, and reacting the resulting compound of formula IV

20

25



30

wherein Z is a leaving group, with pyrrolidine, piperidine, hexamethyleneimine, methylpyrrolidine, dimethylpyrrolidine, or morpholine;


c) deprotecting the 6- and 4'-position hydroxy group of the reaction product of step a) or step b); and
d) optionally salifying or forming a solvate of the reaction product of step c).

35

8. A compound of formula II

40


45



50

9. A compound of formula IV

55



15 wherein Z is a leaving group, particularly bromo, chloro, or iodo.

10. A compound according to Claim 9 wherein said leaving group is bromo.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 95 30 2076

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
A	EP-A-0 062 505 (ELI LILLY AND COMPANY) * the whole document *	1-9	C07D333/56 C07D333/58
D	& US-A-4 380 635 ---		
A	EP-A-0 062 504 (ELI LILLY AND COMPANY) * the whole document *	1-9	
A	GB-A-1 357 212 (LABAZ) * the whole document *	1-9	
A	FR-A-2 329 271 (ELI LILLY AND COMPANY) * the whole document *	1-9	
D	& US-A-4 133 814 ---		
A	US-A-4 024 273 (L. MARTIN BRENNER ET AL.) * the whole document *	1-9	
A	SYNTHESIS, January 1981 pages 1-28, OYO MITSUNOBU 'The use of Diethyl Azodicarboxylate and Triphenylphosphine in Synthesis and Transformation of Natural products' * page 4, column 2, paragraph 2.4 - page 5, column 1 *	1	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 4 August 1995	Examiner Kyriakakou, G
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document	

EPF FORM 1503 (01.92) (P04001)